

Universal behavior in the scattering of heavy, weakly interacting dark matter on nuclear targets

RICHARD J. HILL* AND MIKHAIL P. SOLON†

*Enrico Fermi Institute and Department of Physics
The University of Chicago, Chicago, Illinois, 60637, USA*

Abstract

Particles that are heavy compared to the electroweak scale ($M \gg m_W$), and that are charged under electroweak $SU(2)$ gauge interactions display universal properties such as a characteristic fine structure in the mass spectrum induced by electroweak symmetry breaking, and an approximately universal cross section for scattering on nuclear targets. The heavy particle effective theory framework is developed to compute these properties. As illustration, the spin independent cross section for low-velocity scattering on a nucleon is evaluated in the limit $M \gg m_W$, including complete leading-order matching onto quark and gluon operators, renormalization analysis, and systematic treatment of perturbative and hadronic-input uncertainties.

*richardhill@uchicago.edu

†mpsolon@uchicago.edu

1 Introduction

Cosmological evidence for dark matter consistent with thermal relic Weakly Interacting Massive Particles (WIMPs) motivates laboratory searches for such particles interacting with nuclear targets. Search strategies and detection potential are highly dependent on the WIMP mass, M , and its interaction strength with nuclear matter. We consider here the class of models where the WIMP belongs to an electroweak $SU(2)$ multiplet. This study is motivated in part by the observation that an exact discrete parity arises in Standard Model extensions involving confined fermions coupled to electroweak $SU(2)$ [1]. The parity ensures stability of the lightest pseudo-Nambu Goldstone mode, which is the electrically neutral component of a Lorentz scalar, electroweak $SU(2)$ isotriplet [2].

Regardless of the origin for such an $SU(2)$ multiplet, e.g. whether it is a composite or fundamental particle, universal behavior emerges in the limit where the WIMP mass is large compared to the electroweak scale, $M \gg m_W$. The emergence of these universal properties, and corrections to them, can be systematically analyzed using techniques of heavy particle effective theories [3]. We focus on the case of a real scalar transforming as a triplet of electroweak $SU(2)$, although the results extend straightforwardly to arbitrary $SU(2)$ representations, and to higher spin particles.

At energy scales large compared to m_W , the new particle is described by an effective heavy particle $SU(2)$ gauge theory,

$$\mathcal{L}_{\text{eff}} = \phi_v^* (i v \cdot D + \dots) \phi_v, \quad (1)$$

where ϕ_v is a scalar heavy particle field, v^μ is the heavy particle velocity, and D_μ is the $SU(2)$ covariant derivative. The leading interactions are thus universal, and corrections depending on the mass, or other characteristics of the dark matter particle, are suppressed by powers of M . In this paper we determine the general structure of the heavy scalar effective theory through $\mathcal{O}(1/M^3)$. As an illustrative application, we compute the universal cross section for low-velocity scattering of $SU(2)$ -charged WIMPs on a nucleon in the limit $M \gg m_W$. We present a complete leading order matching onto gluonic operators, renormalization analysis, and systematic treatment of perturbative and hadronic-input uncertainties.

The remainder of the paper is structured as follows. In Section 2, we construct the relevant heavy particle effective theory at scales $\mu \gg m_W$, and compute the leading Wilson coefficients in a simple model. In Section 3 we consider the operator basis and mass corrections in the low energy theory after integrating out scales $\mu \sim m_W$. In Section 4 we perform explicit matching computations at the scale $\mu \sim m_W$. In Section 5 we perform renormalization group evolution from $\mu \sim m_W$ down to low scales $\mu \approx 1 \text{ GeV} \gtrsim \Lambda_{\text{QCD}}$, where QCD matrix elements are estimated. Section 6 presents the cross section for low-velocity scattering on a nucleon. Section 7 presents a summary and outlook.

2 Heavy scalar effective theory: electroweak symmetric theory

Consider the effective theory and matching conditions for a scalar particle of mass M , charged under electroweak $SU(2)$. With obvious modifications, the construction applies to general

gauge groups. We start by investigating the effective theory at scales $m_W \ll \mu \ll M$, with unbroken electroweak gauge symmetry.

2.1 Lagrangian

We work in terms of an effective heavy scalar field $\phi_v(x)$, in the isospin J representation of $SU(2)$. The covariant derivative is $D_\mu = \partial_\mu - ig_2 W_\mu^a t_J^a$ and $W_{\mu\nu} \equiv i[D_\mu, D_\nu]/g_2 \equiv W_{\mu\nu}^a t_J^a$ is the associated field strength. We let $g_1, g_2, g_3 \equiv g$ denote the Standard Model $U(1)_Y, SU(2)_W$ and $SU(3)_c$ gauge coupling constants, respectively. A typical heavy particle momentum can be decomposed as

$$p^\mu = Mv^\mu + k^\mu, \quad (2)$$

where v^μ is a velocity, $v^2 = 1$, and k^μ is a residual momentum. The basis of operators involves the perpendicular derivative,

$$D_\perp^\mu \equiv D^\mu - v^\mu v \cdot D. \quad (3)$$

Through $\mathcal{O}(1/M^3)$, the scalar heavy particle effective theory in the one-heavy-particle sector takes the form,

$$\begin{aligned} \mathcal{L}_\phi = \phi_v^* \bigg\{ & iv \cdot D - c_1 \frac{D_\perp^2}{2M} + c_2 \frac{D_\perp^4}{8M^3} + g_2 c_D \frac{v^\alpha [D_\perp^\beta, W_{\alpha\beta}]}{8M^2} + ig_2 c_M \frac{\{D_\perp^\alpha, [D_\perp^\beta, W_{\alpha\beta}]\}}{16M^3} \\ & + g_2^2 c_{A1} \frac{W^{\alpha\beta} W_{\alpha\beta}}{16M^3} + g_2^2 c_{A2} \frac{v_\alpha v^\beta W^{\mu\alpha} W_{\mu\beta}}{16M^3} + g_2^2 c_{A3} \frac{\text{Tr}(W^{\alpha\beta} W_{\alpha\beta})}{16M^3} + g_2^2 c_{A4} \frac{v_\alpha v^\beta \text{Tr}(W^{\mu\alpha} W_{\mu\beta})}{16M^3} \\ & + g_2^2 c'_{A1} \frac{\epsilon^{\mu\nu\rho\sigma} W_{\mu\nu} W_{\rho\sigma}}{16M^3} + g_2^2 c'_{A2} \frac{\epsilon^{\mu\nu\rho\sigma} v^\alpha v_\mu W_{\nu\alpha} W_{\rho\sigma}}{16M^3} + g_2^2 c'_{A3} \frac{\epsilon^{\mu\nu\rho\sigma} \text{Tr}(W_{\mu\nu} W_{\rho\sigma})}{16M^3} \\ & + g_2^2 c'_{A4} \frac{\epsilon^{\mu\nu\rho\sigma} v^\alpha v_\mu \text{Tr}(W_{\nu\alpha} W_{\rho\sigma})}{16M^3} + \dots \bigg\} \phi_v, \quad (4) \end{aligned}$$

where we have employed appropriate field redefinitions to remove possible redundant operators involving factors of $v \cdot D$ acting on ϕ_v . Note that the operators with coefficients c'_{A1} through c'_{A4} violate parity and time reversal symmetries.¹ For the effective theory describing a fundamental heavy scalar particle, we have $c_1 = c_2 = c_{A1} = 1$ and $c_D = c_M = c_{A2} = c_{A3} = c_{A4} = c'_{A1} = c'_{A2} = c'_{A3} = c'_{A4} = 0$ at tree level [4]. We find that the low-energy manifestation of relativistic invariance (“reparameterization invariance” [5, 6]) implies the exact relations,

$$c_1 = c_2 = 1, \quad c_M = c_D. \quad (5)$$

Section 2.2 provides a nontrivial illustration of the latter relation.

The complete lagrangian including Standard Model particles and interactions can be written

$$\mathcal{L} = \mathcal{L}_\phi + \mathcal{L}_{\text{SM}} + \mathcal{L}_{\phi, \text{SM}}. \quad (6)$$

¹ Additional CPT violating operators at $\mathcal{O}(1/M^2)$ and $\mathcal{O}(1/M^3)$ are constrained by reparameterization invariance to have vanishing coefficient.

Here $\mathcal{L}_{\text{SM}}$ is the usual Standard Model lagrangian, and by convention we have included interactions with W_μ in $\mathcal{L}_\phi$. So far our discussion applies to a general irreducible $SU(2)$ representation for the heavy scalar field ϕ_v . Specializing to the case of a real scalar field, necessarily with integer isospin, the effective theory is invariant under²

$$v_\mu \leftrightarrow -v_\mu, \quad \phi_v \leftrightarrow \phi_v^*. \quad (7)$$

It is straightforward to verify that all interactions in $\mathcal{L}_\phi$ are invariant under this transformation.

In the one-heavy-particle sector, the remaining terms involving the Higgs field H , gauge fields, and fermions are ($\tilde{H} \equiv i\tau_2 H^*$)

$$\begin{aligned} \mathcal{L}_{\phi, \text{SM}} = \phi_v^* \Big\{ & c_H \frac{H^\dagger H}{M} + \dots + c_Q \frac{t_J^a \bar{Q}_L \tau^a \not{Q}_L}{M^2} + c_X \frac{i \bar{Q}_L \tau^a \gamma^\mu Q_L \{t_J^a, D_\mu\}}{2M^3} + c_{DQ} \frac{\bar{Q}_L \not{v} i v \cdot D Q_L}{M^3} \\ & + c_{Du} \frac{\bar{u}_R \not{v} i v \cdot D u_R}{M^3} + c_{Dd} \frac{\bar{d}_R \not{v} i v \cdot D d_R}{M^3} + c_{Hd} \frac{\bar{Q}_L H d_R + h.c.}{M^3} + c_{Hu} \frac{\bar{Q}_L \tilde{H} u_R + h.c.}{M^3} \\ & + g^2 c_{A1}^{(G)} \frac{G^{A\alpha\beta} G_{\alpha\beta}^A}{16M^3} + g^2 c_{A2}^{(G)} \frac{v_\alpha v^\beta G^{A\mu\alpha} G_{\mu\beta}^A}{16M^3} + g^2 c_{A1}'^{(G)} \frac{\epsilon^{\mu\nu\rho\sigma} G_{\mu\nu}^A G_{\rho\sigma}^A}{16M^3} + g^2 c_{A2}'^{(G)} \frac{\epsilon^{\mu\nu\rho\sigma} v^\alpha v_\mu G_{\nu\alpha}^A G_{\rho\sigma}^A}{16M^3} \\ & + \dots \Big\} \phi_v. \end{aligned} \quad (8)$$

Terms odd under (7) have been omitted. Subleading terms containing only H , ϕ_v and their covariant derivatives are represented by the first ellipsis in (8). Terms bilinear in lepton fields, and terms bilinear in the hypercharge gauge field are also present in $\mathcal{L}_{\phi, \text{SM}}$ but have not been written explicitly. Repeated indices $a = 1..3$ and $A = 1..8$ imply a sum over gauge generators. Reparameterization invariance implies

$$c_Q = c_X. \quad (9)$$

2.2 Sample matching calculation

As an illustration of the construction and matching conditions for the heavy particle lagrangian $\mathcal{L}_\phi$, consider the case of a fundamental scalar, ignoring scalar self interactions (i.e., ϕ^4 terms). For the matching of the terms containing a single gauge field, we consider the full theory result for the $W\phi\phi$ amputated three point function,

$$\begin{aligned} & \text{Diagram: A circle with a diagonal line through it, representing a scalar loop. Two external lines (momenta p and p') enter from the top left and top right. A third external line (momentum q) enters from the bottom, represented by a wavy line. The diagram is labeled with p, p', and q and a μ index on the wavy line.} \\ & = i g_2 (p + p')^\mu F(q^2) (t_J^a)_{ji}, \end{aligned} \quad (10)$$

where $q = p' - p$, and $F(q^2)$ is a model-dependent form factor. Setting $p^2 = p'^2 = M^2$, $v^\mu = (1, 0, 0, 0)$, the matching conditions for scalar scattering from a $\mu = 0$ or $\mu = i$ gauge

² For a real scalar field, the effective theory is obtained by introducing v_μ in the field redefinition $\phi(x) = e^{-iMv \cdot x} \phi_v(x) / \sqrt{M} = e^{iMv \cdot x} \phi_v^*(x) / \sqrt{M} = \phi^*(x)$.

field read

$$\begin{aligned}
F(0) - F'(0)\mathbf{q}^2 + \dots &= 1 - c_D \frac{\mathbf{q}^2}{8M^2} + \dots, \\
(p + p')^i \left[-F(0) \left(1 - \frac{\mathbf{p}^2 + \mathbf{p}'^2}{4M^2} \right) + F'(0)\mathbf{q}^2 + \dots \right] \\
&= (p + p')^i \left[-1 + \frac{\mathbf{p}^2 + \mathbf{p}'^2}{4M^2} + c_M \frac{\mathbf{q}^2}{8M^2} \right] + q^i \frac{\mathbf{p}'^2 - \mathbf{p}^2}{8M^2} (c_D - c_M) + \dots
\end{aligned} \tag{11}$$

An explicit computation of one-loop gauge boson corrections, employing dimensional regularization in $d = 4 - 2\epsilon$ dimensions, yields

$$F(q^2) = 1 + \frac{g_2^2}{(4\pi)^2} \frac{q^2}{M^2} \left\{ C_2(r) \left[-\frac{2}{3\epsilon_{\text{IR}}} - 1 + \frac{4}{3} \log \frac{M}{\mu} \right] + C_2(G) \left[-\frac{1}{24\epsilon_{\text{IR}}} + \frac{3}{4} + \frac{1}{12} \log \frac{M}{\mu} \right] \right\} + \dots \tag{12}$$

The quadratic Casimir coefficients for the isospin- J and adjoint representations of $SU(2)$ are $C_2(J) = J(J+1)$ and $C_2(G) = 2$. From (11) and (12), after effective theory subtractions the renormalized coefficients $c_D(\mu)$, $c_M(\mu)$ in the $\overline{\text{MS}}$ renormalization scheme are found to be

$$c_D(\mu) = c_M(\mu) = \frac{\alpha_2(\mu)}{4\pi} \left[-8J(J+1) + 12 + \left(\frac{32J(J+1)}{3} + \frac{4}{3} \right) \log \frac{M}{\mu} \right]. \tag{13}$$

Matching for a general ultraviolet completion model, and for other effective theory coefficients proceeds similarly.

Our focus will be on the limit $M \gg m_W$, where all nontrivial matching conditions at the scale $\mu \sim M$ become irrelevant.³ We leave a detailed investigation of the model-dependent form factor and subleading $1/M$ corrections to future work. Presently we proceed to investigate the leading order predictions of the effective theory at scales $\mu \ll M$.

3 Integrating out the weak scale

For scattering phenomena at $\sim \text{keV}$ energy scales of interest to dark matter-nucleus scattering search experiments, we should examine the appropriate effective theory far below the electroweak scale. Let us begin by integrating out the degrees of freedom at the scale m_W . For definiteness we treat the top quark mass m_t and the Higgs boson mass m_h as parametrically of the same order as m_W . In following sections, we will renormalize to lower energy scales, integrating out the remaining heavy quark degrees of freedom as we pass the bottom and charm quark thresholds. The remaining hadronic matrix elements may then be evaluated in $n_f = 3$ flavor QCD to obtain cross section predictions.

³ In particular models with multiple mass scales, $1/M$ prefactors can be replaced by inverse powers of a smaller excitation energy. It is also of interest to investigate whether large anomalous dimensions could enhance the coefficients of particular subleading operators.

3.1 Mass correction from electroweak symmetry breaking

We may evaluate the heavy scalar self energy to obtain mass corrections,

$$-i\Sigma(p) = \begin{array}{c} W \\ \text{---} \end{array} + \begin{array}{c} Z \\ \text{---} \end{array} + \begin{array}{c} \gamma \\ \text{---} \end{array} + \dots \quad (14)$$

The shift in mass due to electroweak symmetry breaking appears as a nonvanishing value of $\Sigma(p)$ at $v \cdot p = 0$. We find at leading order in the $1/M$ expansion, and first order in perturbation theory,

$$\delta M = \alpha_2 m_W \left[-\frac{1}{2} J^2 + \sin^2 \frac{\theta_W}{2} J_3^2 \right]. \quad (15)$$

In particular, with $Q = J_3 + Y = J_3$ for $Y = 0$, the mass of each charged state is lifted proportional to its squared charge relative to the neutral component,

$$M_{(Q)} - M_{(Q=0)} = \alpha_2 Q^2 m_W \sin^2 \frac{\theta_W}{2} + \mathcal{O}(1/M) \approx (170 \text{ MeV}) Q^2. \quad (16)$$

Subleading corrections can be similarly evaluated in the effective theory. Since no additional operators appear at $\mathcal{O}(1/M^0)$, the result (16) is model independent.⁴

3.2 Operator basis

The effective theory after electroweak symmetry breaking will include: the heavy scalar QED theory for each of the electric charge eigenstates, with mass determined as in (15);⁵ the Standard Model lagrangian with W, Z, h, t integrated out; and interactions,

$$\mathcal{L} = \mathcal{L}_{\phi_0} + \mathcal{L}_{\text{SM}} + \mathcal{L}_{\phi_0, \text{SM}} + \dots, \quad (17)$$

where the ellipsis denotes terms containing electrically charged heavy scalars. For the electrically neutral scalar,

$$\mathcal{L}_{\phi_0} = \phi_{v, Q=0}^* \left\{ i v \cdot \partial - \frac{\partial_\perp^2}{2M_{(Q=0)}} + \mathcal{O}(1/m_W^3) \right\} \phi_{v, Q=0}. \quad (18)$$

Note that enforcing the reality condition (7) implies the vanishing of c_D ($= c_M$).

Interactions with Standard Model fields begin at order $1/m_W^3$. We restrict attention to quark and gluon operators (neglecting lepton and photon operators) and again focus on the neutral $\phi_{v, Q=0}$ component, dropping the $Q = 0$ subscript in the following. Mixing with charged scalars will become relevant at order $1/m_W^4$ in nuclear scattering computations; similarly, we restrict attention to flavor-singlet quark bilinears, since matrix elements of flavor-changing bilinears are suppressed by additional weak coupling factors. Finally, we neglect operators

⁴ The mass splitting (16) appears in limits of particular models, e.g. [1, 7, 8].

⁵ We define the pole mass to include the contributions induced by electroweak symmetry breaking, as opposed to introducing residual mass terms for different charge eigenstates [9].



Figure 1: Matching condition for quark operators. Double lines denote heavy scalars, zigzag lines denote W bosons, dashed lines denote Higgs bosons, single lines with arrows denote quarks, and the solid square denotes an effective theory vertex. Diagrams with crossed W lines are not displayed.

with derivatives acting on ϕ_v or involving γ_5 , since these lead to spin-dependent interactions that are suppressed for low-velocity scattering. The basis of operators is then

$$\mathcal{L}_{\phi_0, \text{SM}} = \frac{1}{m_W^3} \phi_v^* \phi_v \left\{ \sum_q \left[c_{1q}^{(0)} O_{1q}^{(0)} + c_{1q}^{(2)} v_\mu v_\nu O_{1q}^{(2)\mu\nu} \right] + c_2^{(0)} O_2^{(0)} + c_2^{(2)} v_\mu v_\nu O_2^{(2)\mu\nu} \right\} + \dots, \quad (19)$$

where we have chosen QCD operators of definite spin,

$$\begin{aligned} O_{1q}^{(0)} &= m_q \bar{q} q, & O_2^{(0)} &= (G_{\mu\nu}^A)^2, \\ O_{1q}^{(2)\mu\nu} &= \bar{q} \left(\gamma^{\{\mu} i D^{\nu\}} - \frac{1}{d} g^{\mu\nu} i \not{D} \right) q, & O_2^{(2)\mu\nu} &= -G^{A\mu\lambda} G^{A\nu}{}_\lambda + \frac{1}{d} g^{\mu\nu} (G_{\alpha\beta}^A)^2. \end{aligned} \quad (20)$$

Here $A^{\{\mu} B^{\nu\}} \equiv (A^\mu B^\nu + A^\nu B^\mu)/2$ denotes symmetrization. We employ dimensional regularization with $d = 4 - 2\epsilon$ the spacetime dimension. We use the background field method for gluons in the effective theory thus ignoring gauge-variant operators, and assume that appropriate field redefinitions are employed to eliminate operators that vanish by leading order equations of motion. The matrix elements of the gluonic operators, $O_2^{(S)}$, are numerically large, representing a substantial contribution of gluons to the energy and momentum of the nucleon. To account for the leading contributions from both quark and gluon operators, we compute the coefficients $c_2^{(S)}$ through $\mathcal{O}(\alpha_s)$ and $c_{1q}^{(S)}$ through $\mathcal{O}(\alpha_s^0)$.

4 Weak scale matching

The matching conditions for quark operators in the $n_f = 5$ flavor theory at renormalization scale $\mu = \mu_t \sim m_t \sim m_W \sim m_h$ are obtained from the diagrams in Fig. (1):

$$\begin{aligned} c_{1U}^{(0)}(\mu_t) &= \mathcal{C} \left[-\frac{1}{x_h^2} \right], & c_{1D}^{(0)}(\mu_t) &= \mathcal{C} \left[-\frac{1}{x_h^2} - |V_{tD}|^2 \frac{x_t}{4(1+x_t)^3} \right], \\ c_{1U}^{(2)}(\mu_t) &= \mathcal{C} \left[\frac{2}{3} \right], & c_{1D}^{(2)}(\mu_t) &= \mathcal{C} \left[\frac{2}{3} - |V_{tD}|^2 \frac{x_t(3+6x_t+2x_t^2)}{3(1+x_t)^3} \right], \end{aligned} \quad (21)$$

where subscript U denotes u or c and subscript D denotes d , s or b . Here $\mathcal{C} = [\pi\alpha_2^2(\mu_t)][J(J+1)/2]$, $x_h \equiv m_h/m_W$ and $x_t \equiv m_t/m_W$. We ignore corrections of order m_q/m_W for $q = u, d, s, c, b$, and have used CKM unitarity to simplify the results.




Figure 2: Matching condition onto gluon operators. The notation is as in Fig. 1.

Matching conditions onto gluon operators are from the diagrams of Fig. (2):

$$\begin{aligned}
c_2^{(0)}(\mu_t) &= \mathcal{C} \frac{\alpha_s(\mu_t)}{4\pi} \left[\frac{1}{3x_h^2} + \frac{3 + 4x_t + 2x_t^2}{6(1 + x_t)^2} \right], \\
c_2^{(2)}(\mu_t) &= \mathcal{C} \frac{\alpha_s(\mu_t)}{4\pi} \left[-\frac{32}{9} \log \frac{\mu_t}{m_W} - 4 - \frac{4(2 + 3x_t)}{9(1 + x_t)^3} \log \frac{\mu_t}{m_W(1 + x_t)} \right. \\
&\quad - \frac{4(12x_t^5 - 36x_t^4 + 36x_t^3 - 12x_t^2 + 3x_t - 2)}{9(x_t - 1)^3} \log \frac{x_t}{1 + x_t} - \frac{8x_t(-3 + 7x_t^2)}{9(x_t^2 - 1)^3} \log 2 \\
&\quad \left. - \frac{48x_t^6 + 24x_t^5 - 104x_t^4 - 35x_t^3 + 20x_t^2 + 13x_t + 18}{9(x_t^2 - 1)^2(1 + x_t)} \right]. \tag{22}
\end{aligned}$$

There is no dependence of $c_2^{(0)}$ or $c_2^{(2)}$ on CKM matrix elements in the limit of vanishing d, s, b quark masses. The renormalized coefficients are computed in the $\overline{\text{MS}}$ scheme. We have employed Fock-Schwinger ($x \cdot A = 0$) gauge [10] to compute the full-theory amplitudes for gluonic operators in Fig. 2. The effective theory subtractions are efficiently performed in a scheme with massless light quarks, using dimensional regularization as infrared regulator. We have verified that the same results are obtained using finite masses and taking the limit $m_q/m_W \rightarrow 0$. Details of this computation will be presented elsewhere.

5 RG evolution to hadronic scales

To account for perturbative corrections involving large logarithms, e.g. $\alpha_s(\mu_0) \log m_t/\mu_0$, we employ renormalization group evolution to sum leading logarithms to all orders.

5.1 Anomalous dimensions

The spin $S = 0$ and spin $S = 2$ operators mix amongst themselves, with

$$\frac{d}{d \log \mu} O_i^{(S)} = - \sum_j \gamma_{ij}^{(S)} O_j, \quad (23)$$

where $\gamma_{ij}^{(S)}$ are $(n_f + 1) \times (n_f + 1)$ anomalous dimension matrices. The leading terms are

$$\begin{aligned} \hat{\gamma}^{(0)} &= \left(\begin{array}{ccc|ccc} 0 & & & 0 & & \\ & \ddots & & \vdots & & \\ & & 0 & 0 & & \\ \hline -2\gamma'_m & \cdots & -2\gamma'_m & (\beta/g)' & & \end{array} \right) = \frac{\alpha_s}{4\pi} \left(\begin{array}{ccc|ccc} 0 & & & 0 & & \\ & \ddots & & \vdots & & \\ & & 0 & 0 & & \\ \hline 32 & \cdots & 32 & -2\beta_0 & & \end{array} \right) + \dots, \\ \hat{\gamma}^{(2)} &= \frac{\alpha_s}{4\pi} \left(\begin{array}{ccc|ccc} \frac{64}{9} & & & -\frac{4}{3} & & \\ & \ddots & & \vdots & & \\ & & \frac{64}{9} & -\frac{4}{3} & & \\ \hline -\frac{64}{9} & \cdots & -\frac{64}{9} & \frac{4n_f}{3} & & \end{array} \right) + \dots, \end{aligned} \quad (24)$$

where $\beta = dg/d \log \mu \approx -\beta_0 \alpha_s / 4\pi$, $\gamma_m = d \log m_q / d \log \mu \approx -8\alpha_s / 4\pi$, $\gamma'_m \equiv g \partial \gamma_m / \partial g$, $(\beta/g)' \equiv g \partial (\beta/g) / \partial g$, and $\beta_0 = 11 - \frac{2}{3}n_f$. It is straightforward to include subleading terms for $\hat{\gamma}^{(0)}$ [11, 12] and $\hat{\gamma}^{(2)}$ [13, 14].

5.2 Integrating out heavy quarks

At the scale $\mu = \mu_b \sim m_b$, we match onto an $n_f = 4$ theory containing u, d, s, c quarks. The matching equations are

$$\begin{aligned} c_2^{(0)}(\mu_b) &= \tilde{c}_2^{(0)}(\mu_b) \left(1 + \frac{4\tilde{a}}{3} \log \frac{m_b}{\mu_b} \right) - \frac{\tilde{a}}{3} \tilde{c}_{1b}^{(0)}(\mu_b) \left[1 + \tilde{a} \left(11 + \frac{4}{3} \log \frac{m_b}{\mu_b} \right) \right] + \mathcal{O}(\tilde{a}^3), \\ c_{1q}^{(0)}(\mu_b) &= \tilde{c}_{1q}^{(0)}(\mu_b) + \mathcal{O}(\tilde{a}^2), \\ c_2^{(2)}(\mu_b) &= \tilde{c}_2^{(2)}(\mu_b) - \frac{4\tilde{a}}{3} \log \frac{m_b}{\mu_b} \tilde{c}_{1b}^{(2)}(\mu_b) + \mathcal{O}(\tilde{a}^2), \\ c_{1q}^{(2)}(\mu_b) &= \tilde{c}_{1q}^{(2)}(\mu_b) + \mathcal{O}(\tilde{a}), \end{aligned} \quad (25)$$

where $q = u, d, s, c$ and $\tilde{a} = \alpha_s(\mu_b, n_f = 5) / 4\pi$. Quantities without (with) tilde refer to the $n_f = 4$ ($n_f = 5$) theory. The scheme dependence for heavy quark masses enters at higher order. For definiteness we use pole masses for m_b and m_c , with values taken from [15]. Following our power counting scheme, we consider one less order of α_s in the matching for $c_{1q}^{(S)}$ relative to $c_2^{(S)}$. For later use in the numerical analysis, we have included NLO QCD corrections in the spin-0 matching [16, 17]. Similar to above, we evolve coefficients in the $n_f = 4$ theory to the scale $\mu = \mu_c \sim m_c$. Finally, we match onto $n_f = 3$ and evolve to a low scale $\mu_0 \sim 1 \text{ GeV}$ independent of heavy quark masses.

6 Matrix elements and cross section

Having expressed the lagrangian in terms of operators renormalized at the scale $\mu_0 \sim 1 \text{ GeV}$, we require hadronic matrix elements evaluated at this scale.

6.1 Hadronic inputs

Let us define the zero-momentum matrix elements of renormalized operators⁶

$$\begin{aligned}\langle N|O_{1q}^{(0)}|N\rangle &\equiv m_N f_{q,N}^{(0)}, & \frac{-9\alpha_s(\mu)}{8\pi} \langle N|O_2^{(0)}(\mu)|N\rangle &\equiv m_N f_{G,N}^{(0)}(\mu), \\ \langle N|O_{1q}^{(2)\mu\nu}(\mu)|N\rangle &\equiv \frac{1}{m_N} \left(k^\mu k^\nu - \frac{g^{\mu\nu}}{4} m_N^2 \right) f_{q,N}^{(2)}(\mu), \\ \langle N|O_2^{(2)\mu\nu}(\mu)|N\rangle &\equiv \frac{1}{m_N} \left(k^\mu k^\nu - \frac{g^{\mu\nu}}{4} m_N^2 \right) f_{G,N}^{(2)}(\mu),\end{aligned}\tag{26}$$

where m_N is the nucleon mass. Matrix elements refer to a definite (but arbitrary) spin state of the nucleon.

6.1.1 Spin zero

We recall that the spin-0 operator matrix elements are not independent, being linked by the relation [18]

$$m_N = (1 - \gamma_m) \sum_q \langle N|m_q \bar{q}q|N\rangle + \frac{\beta}{2g} \langle N|(G_{\mu\nu}^a)^2|N\rangle, \tag{27}$$

derived from the trace of the QCD energy-momentum tensor. Here $N = p$ or n . Neglecting γ_m , $\mathcal{O}(\alpha_s^2)$ contributions to $\beta(g)$, and power corrections in the above formula, the definitions (26) ensure that $f_{G,N}^{(0)}(\mu) \approx 1 - \sum_{q=u,d,s} f_{q,N}^{(0)}$. Corrections arising from (27) are included in the numerical analysis.

For quark operators, define the scale-independent quantities,

$$\Sigma_{\pi N} = \frac{m_u + m_d}{2} \langle p|(\bar{u}u + \bar{d}d)|p\rangle, \quad \Sigma_0 = \frac{m_u + m_d}{2} \langle p|(\bar{u}u + \bar{d}d - 2\bar{s}s)|p\rangle. \tag{28}$$

In the numerical analysis, we will neglect the small contributions proportional to $|V_{td}|^2$ and $|V_{ts}|^2$, so that $c_{1u}^{(0)} = c_{1d}^{(0)}$. Neglecting also the small contribution [19] $(m_d - m_u) \langle p|(\bar{u}u - \bar{d}d)|p\rangle \sim 2 \text{ MeV}$, and using approximate isospin symmetry, we then require, for $N = p$ or n ,

$$m_N(f_{u,N}^{(0)} + f_{d,N}^{(0)}) \approx \Sigma_{\pi N}, \quad m_N f_{s,N}^{(0)} = \frac{m_s}{m_u + m_d} (\Sigma_{\pi N} - \Sigma_0) = \Sigma_s. \tag{29}$$

⁶ We use nonrelativistic normalization for nucleon states, $\langle N(\mathbf{p})|N(\mathbf{p}')\rangle = (2\pi)^3 \delta^3(\mathbf{p} - \mathbf{p}')$.

Parameter	Value	Reference
$ V_{td} $	~ 0	-
$ V_{ts} $	~ 0	-
$ V_{tb} $	~ 1	-
m_u/m_d	0.49(13)	[20]
m_s/m_d	19.5(2.5)	[20]
$\Sigma_{\pi N}^{\text{lat}}$	0.047(9) GeV	[21]
Σ_s^{lat}	0.050(8) GeV	[22]
$\Sigma_{\pi N}$	0.064(7) GeV	[23]
Σ_0	0.036(7) GeV	[24]
m_W	80.4 GeV	[20]
m_t	172 GeV	[15]
m_b	4.75 GeV	[15]
m_c	1.4 GeV	[15]
m_N	0.94 GeV	-
$\alpha_s(m_Z)$	0.118	[20]
$\alpha_2(m_Z)$	0.0338	[20]

Table 1: Inputs to the numerical analysis.

We consider “traditional” values $\Sigma_{\pi N} = 64 \pm 7 \text{ MeV}$ [23] and $\Sigma_0 = 36 \pm 7 \text{ MeV}$ [24], but investigate also the lattice determinations, $\Sigma_{\pi N}^{\text{lat}} = 47 \pm 9 \text{ MeV}$ [21] and $\Sigma_s^{\text{lat}} = 50 \pm 8 \text{ MeV}$ [22].⁷ We adopt PDG values [20] for light-quark mass ratios. A summary of numerical inputs is presented in Table 1.

6.1.2 Spin two

The matrix elements of spin-two operators can be identified as

$$f_{q,p}^{(2)}(\mu) = \int_0^1 dx x [q(x, \mu) + \bar{q}(x, \mu)], \quad (30)$$

where $q(x, \mu)$ and $\bar{q}(x, \mu)$ are parton distribution functions evaluated at scale μ . Neglecting power corrections, the sum of spin two operators in (20) is the traceless part of the QCD energy momentum tensor, hence independent of scale we have $f_{G,p}^{(2)}(\mu) \approx 1 - \sum_{q=u,d,s} f_{q,p}^{(2)}(\mu)$. Using approximate isospin symmetry we set

$$f_{u,n}^{(2)} = f_{d,p}^{(2)}, \quad f_{d,n}^{(2)} = f_{u,p}^{(2)}, \quad f_{s,n}^{(2)} = f_{s,p}^{(2)}. \quad (31)$$

⁷The latter quantity arises from a naive averaging of $\Sigma_s = 31 \pm 15 \text{ MeV}$ [21] and $\Sigma_s = 59 \pm 10 \text{ MeV}$ [25]. See also [26, 27, 28].

$\mu(\text{GeV})$	$f_{u,p}^{(2)}(\mu)$	$f_{d,p}^{(2)}(\mu)$	$f_{s,p}^{(2)}(\mu)$	$f_{G,p}^{(2)}(\mu)$
1.0	0.404(6)	0.217(4)	0.024(3)	0.36(1)
1.2	0.383(6)	0.208(4)	0.027(2)	0.38(1)
1.4	0.370(5)	0.202(4)	0.030(2)	0.40(1)

Table 2: Operator coefficients derived from MSTW PDF analysis [15] at different values of μ .

Table 2 lists coefficient values for renormalization scales $\mu = 1 \text{ GeV}$, $\mu = 1.2 \text{ GeV}$ and $\mu = 1.4 \text{ GeV}$ determined from the parameterization and analysis of [15].

6.2 Cross section

The low-velocity, spin-independent, cross section for WIMP scattering on a nucleus of mass number A and charge Z may be written

$$\sigma_{A,Z} = \frac{m_r^2}{\pi} |Z\mathcal{M}_p + (A - Z)\mathcal{M}_n|^2 \approx \frac{m_r^2 A^2}{\pi} |\mathcal{M}_p|^2, \quad (32)$$

where $\mathcal{M}_p$ and $\mathcal{M}_n$ are the matrix elements for scattering on a proton or neutron respectively⁸, and $m_r = Mm_N/(M + m_N)$ denotes the reduced mass of the dark-matter nucleus system. As described in Section 6.1, $\mathcal{M}_n \approx \mathcal{M}_p$ up to corrections from numerically small CKM factors and isospin violation in nucleon matrix elements. In the $M \gg m_N$ limit, the cross section scales as A^4 . At finite velocity, a nuclear form factor modifies this behavior [29].

As a numerical benchmark, let us compute the cross section for low-momentum scattering on a nucleon for a heavy real scalar in the isospin representation $J = 1$. Figure 3 displays the result, as a function of the unknown Higgs boson mass. Using Table 1, we consider separately the “traditional” inputs $\Sigma_{\pi N}$ and Σ_0 , as well as recent lattice determinations of $\Sigma_{\pi N}^{\text{lat}}$ and Σ_s^{lat} . For each case, separate bands represent the uncertainty due to neglected perturbative QCD corrections, and due to the hadronic inputs. We estimate the impact of higher order perturbative QCD corrections by varying matching scales $m_W^2/2 \leq \mu_t^2 \leq 2m_t^2$, $m_b^2/2 \leq \mu_b^2 \leq 2m_b^2$, $m_c^2/2 \leq \mu_c^2 \leq 2m_c^2$, $1.0 \text{ GeV} \leq \mu_0 \leq 1.4 \text{ GeV}$, adding the errors in quadrature.

The renormalization group running and heavy quark matching for spin-2 operators are evaluated at LO.⁹ For spin-0 operators, we find a large residual uncertainty at LO from μ_0 , μ_c and μ_b scale variation. The RG running from μ_c to μ_0 from (24) is thus evaluated with NNNLO corrections, including contributions to β/g through $\mathcal{O}(\alpha_s^4)$ and γ_m through $\mathcal{O}(\alpha_s^4)$. Accordingly, the spin-0 gluonic matrix element from (27) is also evaluated at NNNLO,

⁸ Explicitly, $\mathcal{M}_N = m_W^{-3} \langle N | \left(\sum_{q=u,d,s} \left[c_{1q}^{(0)} O_{1q}^{(0)} + c_{1q}^{(2)} v_\mu v_\nu O_{1q}^{(2)\mu\nu} \right] + c_2^{(0)} O_2^{(0)} + c_2^{(2)} v_\mu v_\nu O_2^{(2)\mu\nu} \right) | N \rangle$.

⁹ Up to power corrections and subleading $\mathcal{O}(\alpha_s)$ corrections, our evaluation is equivalent to an evaluation in either the $n_f = 4$ or $n_f = 5$ flavors theories, taking the c - and b -quark momentum fractions of the proton as input. We have verified that these results, with the matrix elements taken from [15], are within our error budget.




Figure 3: Cross section for low-velocity scattering on a nucleon for a heavy real scalar in the isospin $J = 1$ representation of $SU(2)$. The dark shaded region represents the 1σ uncertainty from perturbative QCD, estimated by varying factorization scales. The light shaded region represents the 1σ uncertainty from hadronic inputs.

including contributions to β/g through $\mathcal{O}(\alpha_s^4)$ and γ_m through $\mathcal{O}(\alpha_s^3)$. The residual μ_0 scale variation is insignificant compared to other uncertainties. We perform the RG running and heavy quark matching from μ_t to μ_c at NLO. Hadronic input uncertainties from each source in Table 1 and Table 2 are added in quadrature. We have ignored power corrections appearing at relative order $\alpha_s(m_c)\Lambda_{\text{QCD}}^2/m_c^2$; typical numerical prefactors appearing in the coefficients of the corresponding power-suppressed operators [18] suggest that these effects are small.

Due to a partial cancellation between spin-0 and spin-2 matrix elements, the total cross section and the fractional error depend sensitively on subleading perturbative corrections and on the Higgs mass parameter m_h . We find

$$\sigma_p(m_h = 120 \text{ GeV}) = 0.7 \pm 0.1_{-0.3}^{+0.9} \times 10^{-47} \text{ cm}^2, \quad \sigma_p(m_h = 140 \text{ GeV}) = 2.4 \pm 0.2_{-0.6}^{+1.5} \times 10^{-47} \text{ cm}^2, \quad (33)$$

where the first error is from hadronic inputs, assuming Σ_s^{lat} and $\Sigma_{\pi N}^{\text{lat}}$ from Table 1, and the second error represents the effect of neglected higher order perturbative QCD corrections. For the illustrative value $m_h = 120 \text{ GeV}$, and as a function of the scalar strange-quark matrix element Σ_s , we display the separate contributions of each of the quark and gluon operators in Fig. 4.

7 Summary

We have presented the effective theory for heavy, weakly interacting dark matter candidates charged under electroweak $SU(2)$. Having determined the general form of the effective la-




Figure 4: Breakdown of contributions to the matrix element $\mathcal{M}_p$ using the representative values $m_h = 120$ GeV and $\Sigma_{\pi N}^{\text{lat}} = 47(9)$ MeV. The labels $u^{(S)}$, $d^{(S)}$, $s^{(S)}$ and $g^{(S)}$ refer to spin-S up, down, strange and gluon operator contributions, respectively. The thickness represents the 1σ uncertainty from perturbative QCD. The left-hand vertical band corresponds to the lattice value $\Sigma_s^{\text{lat}} = 50(8)$ MeV and the right-hand vertical band corresponds to the range $\Sigma_s = 366(142)$ MeV deduced from $\Sigma_{\pi N}$ and Σ_0 in Table 1.

grangian (4) through $1/M^3$, we demonstrated matching conditions for subleading operators in a simple model. Using the effective theory, we demonstrated universality of the mass splitting induced by electroweak symmetry breaking, and of the cross section for scattering on nuclear matter. Subleading terms in the $1/M$ expansion can be studied systematically using (4).

Our focus has been on the case of an isotriplet real scalar [1]. For this case, relic abundance estimates [8] indicate that $M \lesssim \text{few TeV}$ in order to not overclose the universe. This mass range, combined with the universal cross section, provides a target for future search experiments.

We have presented a complete matching at first nonvanishing order in α_s , and at leading order in small ratios m_W/M , m_b/m_W and Λ_{QCD}/m_c . We performed renormalization group improvement to sum leading logarithms to all orders. The residual dependence on the high matching scale $\mu_t \sim m_t \sim m_W$ represents uncertainty due to uncalculated higher-order perturbative corrections. Assuming the hadronic input Σ_s^{lat} from Table 1, this scale variation is the largest remaining uncertainty on the cross section; its reduction would require higher loop order calculations.

Our high-scale matching results for quark operators (21) and spin-zero gluon operators agree with $m_W/M \rightarrow 0$ results presented by Hisano et al. [30], under the identification $\mu_t = \mu_b = \mu_c$, i.e., a one-step matching onto the $n_f = 3$ theory.¹⁰ This approach neglects

¹⁰ To make the comparison to the scattering amplitude for a heavy Majorana fermion with $\chi = \chi^c$, we use $\chi = \sqrt{2}e^{-imv \cdot x}(h_v + H_v) = \sqrt{2}e^{imv \cdot x}(h_v^c + H_v^c)$, where h_v and H_v are spinor fields with $(1 - \not{v})h_v = (1 + \not{v})H_v = 0$.

large logarithms appearing in coefficient functions. The effective theory analysis provides a systematically improvable method to resum large logarithms, and to estimate both perturbative and hadronic-input uncertainties. Our results for matching onto spin-two gluon operators are new.

With obvious modifications, our results can be extended to higher $SU(2)$ representations (or $SU(2)$ singlets), fermionic WIMPs, and models with additional low-energy field content beyond the Standard Model. The treatment of QCD corrections presented here can be applied to compute scattering amplitudes in related models, e.g. models involving coupling to hypercharge, or supersymmetric models [29, 31, 32].

Acknowledgements

We acknowledge discussions with Y. Bai, G. Paz and J. Zupan. Work supported by NSF Grant 0855039.

Note added. While this paper was in writing, the preprint [33] appeared, mentioning the invariance (7) for Majorana fermions.

References

- [1] Y. Bai and R. J. Hill, Phys. Rev. D **82**, 111701 (2010) [arXiv:1005.0008 [hep-ph]].
- [2] Other models containing dark matter $SU(2)$ multiplets have been studied in: J. Bagnasco, M. Dine, S. D. Thomas, Phys. Lett. **B320**, 99-104 (1994). [hep-ph/9310290]. M. Cirelli, N. Fornengo and A. Strumia, Nucl. Phys. B **753**, 178 (2006) [arXiv:hep-ph/0512090]. T. Hur, D. W. Jung, P. Ko and J. Y. Lee, Phys. Lett. B **696**, 262 (2011) [arXiv:0709.1218 [hep-ph]]. R. Essig, Phys. Rev. D **78**, 015004 (2008) [arXiv:0710.1668 [hep-ph]]. C. Kilic, T. Okui and R. Sundrum, JHEP **1002**, 018 (2010) [arXiv:0906.0577 [hep-ph]]. M. T. Frandsen, F. Sannino, Phys. Rev. **D81**, 097704 (2010). [arXiv:0911.1570 [hep-ph]]. T. Cohen, J. Kearney, A. Pierce, D. Tucker-Smith, [arXiv:1109.2604 [hep-ph]].
- [3] N. Isgur, M. B. Wise, Phys. Lett. **B232**, 113 (1989). W. E. Caswell, G. P. Lepage, Phys. Lett. **B167**, 437 (1986). E. Eichten, B. R. Hill, Phys. Lett. **B234**, 511 (1990).
- [4] Tree level matching for a heavy colored scalar was considered in: A. H. Hoang and P. Ruiz-Femenia, Phys. Rev. D **73**, 014015 (2006) [arXiv:hep-ph/0511102].
- [5] M. E. Luke, A. V. Manohar, Phys. Lett. **B286**, 348-354 (1992). [hep-ph/9205228].
- [6] A. V. Manohar, Phys. Rev. **D56**, 230-237 (1997). [hep-ph/9701294].
- [7] H. -C. Cheng, B. A. Dobrescu, K. T. Matchev, Nucl. Phys. **B543**, 47-72 (1999). [hep-ph/9811316].
- [8] M. Cirelli, N. Fornengo, A. Strumia, Nucl. Phys. **B753**, 178-194 (2006). [hep-ph/0512090].

- [9] A. F. Falk, M. Neubert, M. E. Luke, Nucl. Phys. **B388**, 363-375 (1992). [hep-ph/9204229].
- [10] V. A. Novikov, M. A. Shifman, A. I. Vainshtein, V. I. Zakharov, Fortsch. Phys. **32**, 585 (1984).
- [11] R. Tarrach, Nucl. Phys. **B196**, 45 (1982).
- [12] B. Grinstein, L. Randall, Phys. Lett. **B217**, 335 (1989).
- [13] A. Vogt, S. Moch, J. A. M. Vermaseren, Nucl. Phys. **B691**, 129-181 (2004). [hep-ph/0404111].
- [14] S. Moch, J. A. M. Vermaseren, A. Vogt, Nucl. Phys. **B688**, 101-134 (2004). [hep-ph/0403192].
- [15] A. D. Martin, W. J. Stirling, R. S. Thorne and G. Watt, Eur. Phys. J. C **63**, 189 (2009) [arXiv:0901.0002 [hep-ph]].
- [16] B. A. Ovrut, H. J. Schnitzer, Phys. Lett. **B110**, 139 (1982).
- [17] T. Inami, T. Kubota, Y. Okada, Z. Phys. C **18**, 69-80 (1983).
- [18] M. A. Shifman, A. I. Vainshtein, V. I. Zakharov, Phys. Lett. **B78**, 443 (1978).
- [19] J. Gasser, H. Leutwyler, Phys. Rept. **87**, 77-169 (1982).
- [20] K. Nakamura *et al.* [Particle Data Group Collaboration], J. Phys. G **G37**, 075021 (2010).
- [21] R. D. Young and A. W. Thomas, Phys. Rev. D **81**, 014503 (2010) [arXiv:0901.3310 [hep-lat]].
- [22] J. Giedt, A. W. Thomas and R. D. Young, Phys. Rev. Lett. **103**, 201802 (2009) [arXiv:0907.4177 [hep-ph]].
- [23] M. M. Pavan, I. I. Strakovsky, R. L. Workman, R. A. Arndt, PiN Newslett. **16**, 110-115 (2002). [hep-ph/0111066].
- [24] B. Borasoy, U. -G. Meissner, Annals Phys. **254**, 192-232 (1997). [hep-ph/9607432].
- [25] D. Toussaint and W. Freeman [MILC Collaboration], Phys. Rev. Lett. **103**, 122002 (2009) [arXiv:0905.2432 [hep-lat]].
- [26] K. Takeda *et al.* [JLQCD Collaboration], Phys. Rev. **D83**, 114506 (2011). [arXiv:1011.1964 [hep-lat]].
- [27] H. Ohki *et al.*, Phys. Rev. D **78**, 054502 (2008) [arXiv:0806.4744 [hep-lat]].
- [28] S. Durr, Z. Fodor, T. Hemmert, C. Hoelbling, J. Frison, S. D. Katz, S. Krieg, T. Kurth *et al.*, “Sigma term and strangeness content of octet baryons,” arXiv:1109.4265 [hep-lat].

- [29] G. Jungman, M. Kamionkowski and K. Griest, Phys. Rept. **267**, 195 (1996) [arXiv:hep-ph/9506380].
- [30] J. Hisano, K. Ishiwata, N. Nagata, T. Takesako, JHEP **1107**, 005 (2011). [arXiv:1104.0228 [hep-ph]]. J. Hisano, K. Ishiwata, N. Nagata, Phys. Rev. **D82**, 115007 (2010). [arXiv:1007.2601 [hep-ph]].
- [31] M. Drees, M. M. Nojiri, Phys. Rev. **D47**, 4226-4232 (1993). [hep-ph/9210272]. M. Drees, M. Nojiri, Phys. Rev. **D48**, 3483-3501 (1993). [hep-ph/9307208].
- [32] J. R. Ellis, K. A. Olive and C. Savage, Phys. Rev. D **77**, 065026 (2008) [arXiv:0801.3656 [hep-ph]].
- [33] K. Kopp, T. Okui, [arXiv:1108.2702 [hep-ph]].